arXiv:1404.2245v1 [math.DG] 8 Apr 2014

OPTIMAL GEOMETRIC ESTIMATES FOR FRACTIONAL SOBOLEV CAPACITIES

JIE XIAO

Abstract. This note develops certain sharp inequalities relating the fractional Sobolev capacity of a set to its standard volume and fractional perimeter.

Partially motivated by [20, 21], this note discovers some optimal estimates linking the fractional Sobolev capacity of a set to its standard volume and fractional perimeter.

1. Fractional Sobolev capacities and their basic properties. Let $0 < \alpha < 1$ and C_0^∞ denote the class of all smooth functions with compact support in $\mathbb{R}^n$. Define the fractional Sobolev space (or the homogeneous $(\alpha, 1, 1)$-Besov space) $\dot\Lambda_\alpha^{1,1}$ as the completion of all functions $f \in C_0^\infty$ with

$$\|f\|_{\dot\Lambda_\alpha^{1,1}} = \int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} |f(x+h) - f(x)| dx \right) \frac{dh}{|h|^{n+\alpha}}.$$

Attached to $\dot\Lambda_\alpha^{1,1}$ is the following set-function:

$$\mathrm{cap}(K; \dot\Lambda_\alpha^{1,1}) = \inf \{ \|f\|_{\dot\Lambda_\alpha^{1,1}} : \quad f \in C_0^\infty \quad \& \quad f \ge 1_K \} \quad \forall \quad \text{compact} \quad K \subset \mathbb{R}^n.$$

Here and henceforth, 1_E stands for the indicator of a set $E \subset \mathbb{R}^n$. This definition is extended to any set $E \subset \mathbb{R}^n$ via

$$\mathrm{cap}(E; \dot\Lambda_\alpha^{1,1}) = \inf_{\text{open } O \supseteq E} \mathrm{cap}(O; \dot\Lambda_\alpha^{1,1}) = \inf_{\text{open } O \supseteq E} \left(\sup_{\text{compact } K \subseteq O} \mathrm{cap}(K; \dot\Lambda_\alpha^{1,1}) \right).$$

The number $\mathrm{cap}(E; \dot\Lambda_\alpha^{1,1})$ is called the fractional Sobolev capacity (or the homogeneous end-point Besov capacity) of E; see also [1, 2, 3, 20, 18]. Note that (cf. [16, 17, 4, 5, 12])

$$\lim_{\alpha \to 0} \alpha \|f\|_{\dot\Lambda_\alpha^{1,1}} = 2n\omega_n \|f\|_{L^1} \quad \& \quad \lim_{\alpha \to 1} (1-\alpha) \|f\|_{\dot\Lambda_\alpha^{1,1}} = \tau_n \|\nabla f\|_{L^1}$$

where ω_n is the volume of the unit ball $\mathbb{B}^n$ of $\mathbb{R}^n$ and

$$\tau_n = \int_{\mathbb{S}^{n-1}} |\cos\theta| \, d\sigma$$

with: $\mathbb{S}^{n-1}$ being the unit sphere of $\mathbb{R}^n$; θ being the angle deviation from the vertical direction; and $d\sigma$ being the standard area measure on $\mathbb{S}^{n-1}$. So, we have that for any compact $K \subset \mathbb{R}^n$,

$$\lim_{\alpha \to 0} \alpha \mathrm{cap}(K; \dot\Lambda_\alpha^{1,1}) = 2n\omega_n V(K) \quad \& \quad \lim_{\alpha \to 1} (1-\alpha) \mathrm{cap}(K; \dot\Lambda_\alpha^{1,1}) = \tau_n \mathrm{cap}(K; \dot W^{1,1})$$

where

$$V(K) = \int_K dx \quad \& \quad \mathrm{cap}(K; \dot W^{1,1}) = \inf \{ \|\nabla f\|_{L^1} : \quad f \in C_0^\infty \quad \& \quad f \ge 1_K \}$$

2010 *Mathematics Subject Classification.* Primary 26D10, 31B15, 46E35; Secondary 52A38, 53A15, 53A30.
Key words and phrases. Sharpness, analytic/geometric inequality, volume, fractional capacity/perimeter.
Research supported in part by: NSERC and URP of MUN, Canada.

and

$$\mathrm{cap}(E;\dot W^{1,1})=\inf_{\text{open } O\supseteq E}\mathrm{cap}(O;\dot W^{1,1})=\inf_{\text{open } O\supseteq E}\left(\sup_{\text{compact } K\subseteq O}\mathrm{cap}(K;\dot W^{1,1})\right)\quad\forall\quad E\subset\mathbb{R}^n.$$

Theorem 1. *The nonnegative set-function $E\mapsto cap(E;\dot\Lambda_\alpha^{1,1})$ enjoys three essential properties below:*

(i) *Homogeneity:* $cap(rE;\dot\Lambda_\alpha^{1,1})=r^{n-\alpha}cap(E;\dot\Lambda_\alpha^{1,1})\ \forall\ rE=\{rx:\ x\in E\}\subset\mathbb{R}^n$ & $r\in[0,\infty)$.

(ii) *Monotonicity:* $E_1\subseteq E_2\subset\mathbb{R}^n\Rightarrow cap(E_1;\dot\Lambda_\alpha^{1,1})\le cap(E_2;\dot\Lambda_\alpha^{1,1})$.

(iii) *Upper-semi-continuity:* $\lim_{j\to\infty}cap(K_j;\dot\Lambda_\alpha^{1,1})=cap(\cap_{j=1}^\infty K_j;\dot\Lambda_\alpha^{1,1})\ \forall$ *sequence* $\{K_j\}_{j=1}^\infty$ *of compact subsets of* $\mathbb{R}^n$ *with* $K_1\supseteq K_2\supseteq K_3\supseteq\cdots$.

Proof. (i) follows from

$$\|f(r\cdot)\|_{\dot\Lambda_\alpha^{1,1}}=r^{\alpha-n}\|f\|_{\dot\Lambda_\alpha^{1,1}}\quad\forall\quad r\in[0,\infty).$$

and the definition of $\mathrm{cap}(\cdot,\dot\Lambda_\alpha^{1,1})$.

(ii) follows from the definition of $\mathrm{cap}(\cdot,\dot\Lambda_\alpha^{1,1})$.

(iii) follows from a careful treatment. Suppose $\{K_j\}_{j=1}^\infty$ is a decreasing sequence of compact subsets of $\mathbb{R}^n$. Then $K=\cap_{j=1}^\infty K_j$ is compact. Following the argument for [10, Theorem 2.2(iv)], for any $\epsilon\in(0,1)$ there is a function $f\in C_0^\infty$ such that

$$f\ge1_K\quad\&\quad\|f\|_{\dot\Lambda_\alpha^{1,1}}<\mathrm{cap}(K;\dot\Lambda_\alpha^{1,1})+\epsilon.$$

Note that if j is sufficiently large then K_j is contained in the compact set $\{x\in\mathbb{R}^n:\ f(x)\ge1-\epsilon\}$. So, an application of (ii) and the definition of $\mathrm{cap}(\cdot;\dot\Lambda_\alpha^{1,1})$ derives

$$\lim_{j\to\infty}\mathrm{cap}(K_j;\dot\Lambda_\alpha^{1,1})\le\mathrm{cap}(\{x\in\mathbb{R}^n:\ f(x)\ge1-\epsilon\};\dot\Lambda_\alpha^{1,1})\le\frac{\|f\|_{\dot\Lambda_\alpha^{1,1}}}{1-\epsilon}\le\frac{\mathrm{cap}(K;\dot\Lambda_\alpha^{1,1})+\epsilon}{1-\epsilon}.$$

Upon letting $\epsilon\to0$ and using (ii) again, we get

$$\mathrm{cap}(K;\dot\Lambda_\alpha^{1,1})\le\lim_{j\to\infty}\mathrm{cap}(K_j;\dot\Lambda_\alpha^{1,1})\le\mathrm{cap}(K;\dot\Lambda_\alpha^{1,1}),$$

as desired. □

For any set $E\subset\mathbb{R}^n$, let $E^c=\mathbb{R}^n\setminus E$ and compute

$$\|1_E\|_{\dot\Lambda_\alpha^{1,1}}=2\int_E\int_{E^c}\frac{dxdy}{|x-y|^{n+\alpha}}\equiv2P_\alpha(E).$$

whose half $P_\alpha(E)$ is called the fractional α-perimeter; see e.g. [9, 13]. Notice that

$$\lim_{\alpha\to0}\alpha P_\alpha(E)=n\omega_nV(E)\quad\&\quad\lim_{\alpha\to1}(1-\alpha)P_\alpha(E)=2^{-1}\tau_nP(E)$$

where $P(E)$ is the perimeter of E. So, we get an extension of [15, Lemma 2.2.5] from the limit $\alpha\to1$ to the intermediate value $0<\alpha<1$ that connects the fractional Sobolev capacity and the fractional perimeter.

Theorem 2. *If K is a compact subset of $\mathbb{R}^n$, then*

$$cap(K;\dot\Lambda_\alpha^{1,1})=2\inf_{O\in\mathsf{O}^\infty(K)}P_\alpha(O)$$

where $\mathsf{O}^\infty(K)$ denotes the class of all open sets with C^∞ boundary that contain K.

Proof. Given a compact $K \subset \mathbb{R}^n$. On the one hand, if $f \in C_0^\infty$ and $f \geq 1_K$ then

$$K \subset \{x \in \mathbb{R}^n : f(x) > t\} \quad \forall \quad t \in (0,1),$$

and hence an application of the generalized co-area formula in [19] (cf. [20, Theorem 1.2] for another version of the co-area formula of dimension $n-\alpha$) gives

$$\|f\|_{\dot{\Lambda}_\alpha^{1,1}} = 2\int_0^\infty P_\alpha(\{x \in \mathbb{R}^n : f(x) > t\})\,dt \geq 2\int_0^1 P_\alpha(\{x \in \mathbb{R}^n : f(x) > t\})\,dt \geq 2 \inf_{O \in \mathsf{O}^\infty(K)} P_\alpha(O).$$

This, along with the definition of $\mathrm{cap}(K;\dot{\Lambda}_\alpha^{1,1})$, implies

$$\mathrm{cap}(K;\dot{\Lambda}_\alpha^{1,1}) \geq 2 \inf_{O \in \mathsf{O}^\infty(K)} P_\alpha(O).$$

On the other hand, according to Theorem 1(ii) and [11, Theorem 3.1] we have

$$\mathrm{cap}(K;\dot{\Lambda}_\alpha^{1,1}) \leq \mathrm{cap}(\overline{O};\dot{\Lambda}_\alpha^{1,1}) \leq 2P_\alpha(O) \quad \forall \quad O \in \mathsf{O}^\infty(K),$$

whence

$$\mathrm{cap}(K;\dot{\Lambda}_\alpha^{1,1}) \leq 2 \inf_{O \in \mathsf{O}^\infty(K)} P_\alpha(O).$$

Therefore, the desired formula for $\mathrm{cap}(K;\dot{\Lambda}_\alpha^{1,1})$ follows. □

As an immediate consequence of Theorem 2, we have

$$\lim_{\alpha\to 0} \alpha \mathrm{cap}(K;\dot{\Lambda}_\alpha^{1,1}) = 2n\omega_n V(K) \quad \& \quad \lim_{\alpha\to 1}(1-\alpha)\mathrm{cap}(K;\dot{\Lambda}_\alpha^{1,1}) = \tau_n P(K) \quad \forall \text{ compact } K \subset \mathbb{R}^n.$$

2. Fractional Sobolev inequalities and their geometric forms. The next analytic-geometric assertion indicates that the fractional Sobolev capacity plays a decisive role in improving the fractional isoperimetric inequality [8, (4.2)].

Theorem 3. *Let* $\kappa_{n,\alpha} = \omega_n^{\frac{n-\alpha}{n}}(2P_\alpha(\mathbb{B}^n))^{-1}$. *Then:*
(i) *The analytic inequality*

$$\|f\|_{L^{\frac{n}{n-\alpha}}} \leq \kappa_{n,\alpha}\left(\int_0^\infty (cap(\{x \in \mathbb{R}^n : |f(x)| \geq t\};\dot{\Lambda}_\alpha^{1,1}))^{\frac{n}{n-\alpha}}\,dt^{\frac{n}{n-\alpha}}\right)^{\frac{n-\alpha}{n}} \quad \forall\ f \in C_0^\infty \tag{1}$$

is equivalent to the geometric inequality

$$(V(O))^{\frac{n-\alpha}{n}} \leq \kappa_{n,\alpha}\, cap(\overline{O};\dot{\Lambda}_\alpha^{1,1}) \quad \forall \text{ bounded domain } O \subset \mathbb{R}^n \text{ with } C^\infty \text{ boundary } \partial O. \tag{2}$$

Moreover, both (1) *and* (2) *are true and sharp.*
(ii) *The analytic inequality*

$$\left(\int_0^\infty (cap(\{x \in \mathbb{R}^n : |f(x)| \geq t\};\dot{\Lambda}_\alpha^{1,1}))^{\frac{n}{n-\alpha}}\,dt^{\frac{n}{n-\alpha}}\right)^{\frac{n-\alpha}{n}} \leq \|f\|_{\dot{\Lambda}_\alpha^{1,1}} \ \forall\ f \in C_0^\infty \tag{3}$$

is equivalent to the geometric inequality

$$cap(\overline{O};\dot{\Lambda}_\alpha^{1,1}) \leq 2P_\alpha(O) \quad \forall \text{ bounded domain } O \subset \mathbb{R}^n \text{ with } C^\infty \text{ boundary } \partial O. \tag{4}$$

Moreover, both (3) and (4) are true and sharp.

Proof. (i) Suppose (2) is valid. For any C_0^∞ function f, set

$$O_t(f) = \{x \in \mathbb{R}^n : \ |f(x)| > t\} \quad \forall \quad t \ge 0.$$

Then an application of (2) to $O_t(f)$ yields

$$\|f\|_{\frac{n}{n-s}} = \left(\int_0^\infty V(O_t(f))\, dt^{\frac{n}{n-\alpha}}\right)^{\frac{n-\alpha}{n}} \le \kappa_{n,\alpha}\left(\int_0^\infty \left(\mathrm{cap}(\overline{O_t(f)}; \dot\Lambda_\alpha^{1,1})\right)^{\frac{n}{n-\alpha}} dt^{\frac{n}{n-\alpha}}\right)^{\frac{n-\alpha}{n}},$$

deriving (1). Conversely, suppose (1) is valid. For any bounded domain $O \subset \mathbb{R}^n$ with C^∞ boundary ∂O, the Euclidean distance $\mathrm{dist}(x, E)$ of a point x to a set E, and $0 < \epsilon < 1$, let

$$f_\epsilon(x) = \begin{cases} 1 - \epsilon^{-1}\mathrm{dist}(x, \overline{O}) & \text{as} \quad \mathrm{dist}(x, \overline{O}) < \epsilon \\ 0 \quad \text{as} \quad \mathrm{dist}(x, \overline{O}) \ge \epsilon. \end{cases}$$

Then the inequality in (1) is true for f_ϵ. Consequently, via setting

$$O_\epsilon = \{x \in \mathbb{R}^n : \ \mathrm{dist}(x, \overline{O}) < \epsilon\} \quad \& \quad \epsilon \to 0$$

and using Theorem 1(iii), we gain

$$\begin{aligned} &(V(O))^{\frac{n-\alpha}{n}} \\ &\leftarrow \|f_\epsilon\|_{L^{\frac{n}{n-\alpha}}} \\ &\le \kappa_{n,\alpha}\left(\int_0^\infty (\mathrm{cap}(\overline{O_t(f_\epsilon)}; \dot\Lambda_\alpha^{1,1}))^{\frac{n}{n-\alpha}}\, dt^{\frac{n}{n-\alpha}}\right)^{\frac{n-\alpha}{n}} \\ &= \kappa_{n,\alpha}\left(\int_0^1 (\mathrm{cap}(\overline{O_t(f_\epsilon)}; \dot\Lambda_\alpha^{1,1}))^{\frac{n}{n-\alpha}}\, dt^{\frac{n}{n-\alpha}}\right)^{\frac{n-\alpha}{n}} \\ &\le \kappa_{n,\alpha}\mathrm{cap}(\overline{O_\epsilon}; \dot\Lambda_\alpha^{1,1}) \\ &\to \kappa_{n,\alpha}\mathrm{cap}(\overline{O}; \dot\Lambda_\alpha^{1,1}). \end{aligned}$$

This proves (2). Moreover, the truth and the sharpness of (2) (and hence (1) via the just-checked equivalence) follow from the definition of $\mathrm{cap}(O; \dot\Lambda_\alpha^{1,1})$ and the sharp fractional Sobolev inequality on [8, Theorem 4.1: $p = 1$]:

$$\|f\|_{L^{\frac{n}{n-\alpha}}} \le \kappa_{n,\alpha}\|f\|_{\dot\Lambda_\alpha^{1,1}} \quad \forall \ \ f \in C_0^\infty.$$

(ii) Suppose (3) is valid. Given $\epsilon \in (0, 1)$ and a bounded domain $O \subset \mathbb{R}^n$ with C^∞ boundary ∂O, select again O_ϵ & f_ϵ as above. Then Theorem 1(ii) and the argument for [11, Lemma 3.2] give

$$\mathrm{cap}(\overline{O}; \dot\Lambda_\alpha^{1,1}) \le \left(\int_0^1 \left(\mathrm{cap}(\overline{O_t(f_\epsilon)}; \dot\Lambda_\alpha^{1,1})\right)^{\frac{n}{n-\alpha}} dt^{\frac{n}{n-\alpha}}\right)^{\frac{n-\alpha}{n}} \le \|f_\epsilon\|_{\dot\Lambda_\alpha^{1,1}} \to \|1_O\|_{\dot\Lambda_\alpha^{1,1}} = 2P_\alpha(O).$$

In other words, (4) is true. Conversely, suppose (4) is valid. Upon noticing that for any C_0^∞ function f with $O_t(f)$ being as above, the function $t \mapsto \mathrm{cap}(\overline{O_t(f)}; \ \dot\Lambda_\alpha^{1,1})$ decreases on $[0, \infty)$ (thanks to Theorem 1(ii)), we have

$$\begin{aligned} &t^{\frac{\alpha}{n-\alpha}}\left(\mathrm{cap}(\overline{O_t(f)}; \dot\Lambda_\alpha^{1,1})\right)^{\frac{n}{n-\alpha}} \\ &= \left(t\mathrm{cap}(\overline{O_t(f)}; \ \dot\Lambda_\alpha^{1,1})\right)^{\frac{\alpha}{n-\alpha}}\mathrm{cap}(\overline{O_t(f)}; \ \dot\Lambda_\alpha^{1,1}) \\ &\le \left(\int_0^t \mathrm{cap}(\overline{O_s(f)}; \ \dot\Lambda_\alpha^{1,1})\, ds\right)^{\frac{\alpha}{n-\alpha}}\mathrm{cap}(\overline{O_t(f)}; \ \dot\Lambda_\alpha^{1,1}) \end{aligned}$$

$$= \Big(\frac{n-\alpha}{n}\Big)\frac{d}{dt}\left(\int_0^t \mathrm{cap}(\overline{O_s(f)};\ \dot\Lambda_\alpha^{1,1})\,ds\right)^{\frac{n}{n-\alpha}},$$

whence finding, along with Theorem 1(ii), (4), Theorem 2 and the previously-cited co-area formula,

$$\left(\int_0^\infty \Big(\mathrm{cap}(\overline{O_t(f)});\ \dot\Lambda_\alpha^{1,1}\Big)^{\frac{n}{n-\alpha}}\,dt^{\frac{n}{n-\alpha}}\right)^{\frac{n-\alpha}{n}} \le \int_0^\infty \mathrm{cap}(\overline{O_t(f)};\dot\Lambda_\alpha^{1,1})\,dt \le 2\int_0^\infty P_\alpha(O_t(f))\,dt = \|f\|_{\dot\Lambda_\alpha^{1,1}}.$$

So, (3) holds. Moreover, the truth of (4) (and hence (3) via the above equivalence) follows from Theorem 2. In fact, if (4) were not sharp, then an application of (2) would derive that the sharp fractional isoperimetric inequality (cf. [8, (4.2)])

$$(V(E))^{\frac{n-\alpha}{n}} \le 2\kappa_{n,\alpha}P_\alpha(E) \quad \forall \quad E\subset\mathbb{R}^n$$

is not sharp, thereby reaching a contradiction. Thus, (4) is sharp, and so is (3). □

Theorem 3 comes actually from splitting both the sharp fractional Sobolev inequality and the fractional isoperimetric inequality whose equivalence (optimizing [11, Theorem 1.1] under $G = \mathbb{R}^n$) is described below.

Theorem 4. *The following three optimal statements are equivalent:*

(i) *The fractional Sobolev inequality* $\|f\|_{L^{\frac{n}{n-\alpha}}} \le \kappa_{n,\alpha}\|f\|_{\dot\Lambda_\alpha^{1,1}}$ *holds for any* $f\in C_0^\infty$.

(ii) *The fractional isocapacitary inequality* $(V(O))^{\frac{n-\alpha}{n}} \le \kappa_{n,\alpha}cap(\overline{O},\dot\Lambda_\alpha^{1,1})$ *holds for any bounded domain* $O\subset\mathbb{R}^n$ *with* C^∞ *boundary* ∂O.

(iii) *The fractional isoperimetric inequality* $(V(O))^{\frac{n-\alpha}{n}} \le 2\kappa_{n,\alpha}P_\alpha(O)$ *holds for any bounded domain* $O\subset\mathbb{R}^n$ *with* C^∞ *boundary* ∂O.

Proof. (i)⇒(ii) follows from the definition of $\mathrm{cap}(\overline{O},\dot\Lambda_\alpha^{1,1})$. (ii)⇒(iii) follows from Theorem 2. (iii)⇒(i) follows from the idea verifying (4)⇒(3). As a matter of fact, assume (iii) is true. Given a function $f\in C_0^\infty$ with $O_t(f)$ being the same as in the proof of Theorem 3. Obviously, $t\mapsto V(O_t(f))$ is a decreasing function on $[0,\infty)$. This monotonicity, together with the layer-cake formula, the chain rule, (iii) for $O_t(f)$, and the above-used co-area formula, derives

$$\begin{aligned}
&\|f\|_{L^{\frac{n}{n-\alpha}}}\\
&= \left(\int_0^\infty V(O_t(f))\,dt^{\frac{n}{n-\alpha}}\right)^{\frac{n-\alpha}{n}}\\
&= \int_0^\infty \frac{d}{dt}\left(\int_0^t V(O_s(f))\,ds^{\frac{n}{n-\alpha}}\right)^{\frac{n-\alpha}{n}}dt\\
&= \int_0^\infty \left(\int_0^t V(O_s(f))\,ds^{\frac{n}{n-\alpha}}\right)^{-\frac{\alpha}{n}} V(O_t(f))t^{\frac{\alpha}{n-\alpha}}\,dt\\
&\le \int_0^\infty \Big(V(O_t(f))\Big)^{\frac{n-\alpha}{n}}\,dt\\
&\le 2\kappa_{n,\alpha}\int_0^\infty P_\alpha(O_t(f))\,dt\\
&= \kappa_{n,\alpha}\|f\|_{\dot\Lambda_\alpha^{1,1}},
\end{aligned}$$

whence reaching (i).

□

Note that for any compact set $K \subset \mathbb{R}^n$ and any bounded domain $O \subset \mathbb{R}^n$ one has

$$\begin{cases} \lim_{\alpha\to 1}(1-\alpha)^{-1}\kappa_{n,\alpha} = (n\omega_n^{\frac{1}{n}}\tau_n)^{-1} \\ \lim_{\alpha\to 1}(1-\alpha)\mathrm{cap}(K;\ \dot{\Lambda}_\alpha^{1,1}) = \tau_n \mathrm{cap}(K;\ \dot{W}^{1,1}) = \tau_n \inf_{\tilde{O}\in \mathsf{O}^\infty(K)} P(\tilde{O}) \\ \lim_{\alpha\to 1}(1-\alpha)P_\alpha(O) = 2^{-1}\tau_n P(O). \end{cases}$$

So, the limiting cases of Theorem 3 and Theorem 4 as $\alpha \to 1$ reduce to [21, Theorems 1.1-1.2] and [22, Proposition 3.1] plus the well-known Federer-Felming-Maz'ya equivalence between the isoperimetric inequality and the Sobolev inequality (cf. [7, 14]), respectively.

References

1. D. R. Adams, *Lectures on L^p-potential Theory,* Univ. of Umea, Dept. of Math. 2, 1981.
2. D. R. Adams, Besov capacity redux, *J. Math. Sci.(N.Y.)* 162(2009)307318.
3. D. R. Adams and J. Xiao, Strong type estimates for homogeneous Besov capacities, *Math. Ann.* 325(2003)695-709.
4. J. Bourgain, H. Brezis and P. Mironescu, Another look at Sobolev spaces, in Optimal Control and Partial Differential Equations (J. L. Menaldi, E. Rofman and A. Sulem, eds.) a volume in honour of A. Bensoussan's 60th birthday, IOS Press, 2001, pp. 439-455.
5. J. Bourgain, H. Brezis and P. Mironescu, Limiting embedding theorems for $W^{s,p}$ when $s \uparrow 1$ and applications, *J. Anal. Math.* 87(2002)77-101.
6. J. Dávila, On an open question about functions of bounded variation, *Calc. Var. Partial Differential Equations* 15(2002)519-527.
7. H. Federer and W. H. Fleming, Normal and integral currents, *Ann. Math.* 72(1960)458-520.
8. R. L. Frank and R. Seiringer, Non-linear ground state representations and sharp Hardy inequalities, *J. Funct. Anal.* 252(2008)3407-3430.
9. N. Fusco, V. Millot and M. Morini, A quantitative isoperimetric inequality for fractional perimeters, *J. Funct. Anal.* 261(2011)697-715.
10. J. Heinonen, T. Kilpeläinen, and O. Martio, *Nonlinear Potential Theory of Degenerate Elliptic Equations*, Oxford University Press, 1993.
11. R. Hurri-Syrjänen and A. V. Vähäkangas, Characterizations to the fractional Sobolev inequality, *arXiv:1312.3135v1[math.CA] 11Dec 2013*.
12. M. Ludwig, Anisotropic fractional Sobolev norms, *Adv. Math.* 252(2014)150-157.
13. M. Ludwig, Anisotropic fractional perimeters, *J. Differential Geom.* 96(2014)77-93.
14. V. Maz'ya, Classes of domains and imbedding theorems for function spaces (Russian). Dokl. Akad. Nauk SSSR 3(1960)527-530; English transl.: Sov. Math. Dokl 1(1961)882-885.
15. V. Maz'ya, *Sobolev Spaces with Applications to Elliptic Partial Differential Equations*, 2nd, revised and augmented edition, Springer, 2011.
16. V. Maz'ya and T. Shaposhnikova, On the Bourgain, Brezis, and Mironescu theorem concerning limiting embeddings of fractional Sobolev spaces, *J. Funct. Anal.* 195(2002)230-238.
17. V. Maz'ya and T. Shaposhnikova, Erratum to: "On the Bourgain, Brezis, and Mironescu theorem concerning limiting embeddings of fractional Sobolev spaces" [J. Funct. Anal. 195(2002)230-238], *J. Funct. Anal.* 201(2003)298-300.
18. P. Silvestre, Capacities and embeddings via symmetrization and conductor inequalities, *Proc. Amer. Math. Soc.* 142(2014)497-505.
19. A. Visintin, Nonconvex functionals related to multiphase systems, *SIAM J. Math. Anal.* 21(1990)1281-1304.
20. J. Xiao, Homogeneous endpoint Besov space embeddings by Hausdorff capacity and heat equation, *Adv. Math.* 207(2006)828-846.
21. J. Xiao, The sharp Sobolev and isoperimetric inequalities split twice, *Adv. Math.* 211(2007)417-435.
22. J. Xiao, The p-Faber-Krahn inequality noted, In: Around the Research of Vladimir Maz'ya I. Function Spaces, pp. 373-390, Springer, 2010.

Department of Mathematics and Statistics, Memorial University, St. John's, NL A1C 5S7, Canada
E-mail address: jxiao@mun.ca